

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

Pr **TOBREX® 0.3%** **Tobramycin Ophthalmic Solution**

Read this carefully before you start taking **TOBREX®** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **TOBREX**.

What is TOBREX used for?

TOBREX is used in adults and children above the age of 1 year to treat bacterial infections of the external part of the eye, such as conjunctivitis.

Antibacterial drugs like **TOBREX** treat only bacterial infections. They do not treat viral infections. Although you may feel better early in treatment, **TOBREX** should be used exactly as directed. Misuse or overuse of **TOBREX** could lead to the growth of bacteria that will not be killed by **TOBREX** (resistance). This means that **TOBREX** may not work for you in the future. Do not share your medicine.

How does TOBREX work?

TOBREX contains a medicine called tobramycin. Tobramycin belongs to the aminoglycoside class of antibiotic. Tobramycin works by killing bacteria that cause infections.

The signs of your eye infections [eye redness, increased tearing, discharge (pus or mucus) etc.] may start to improve within 3 days after starting **TOBREX**.

What are the ingredients in TOBREX?

Medicinal ingredients: tobramycin 0.3% w/v (3 mg/mL)

Non-medicinal ingredients:

Preservative: benzalkonium chloride 0.01% w/v

Other ingredients: boric acid, sodium sulfate, sodium chloride, tyloxapol, sulfuric acid and/or sodium hydroxide (to adjust pH) and purified water.

TOBREX comes in the following dosage forms:

TOBREX comes as eye drop and eye ointment.

Do not use TOBREX if you are allergic to:

- tobramycin,
- any of the other ingredients in **TOBREX** (see **What are the ingredients in TOBREX?**),
- other aminoglycoside antibiotics such as amikacin, gentamycin, kanamycin, paromomycin, streptomycin.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take TOBREX. Talk about any health conditions or problems you may have, including if you:

- are pregnant, you think you may be pregnant or are planning to have a baby. You will need to talk to your healthcare professional about the benefits and risks of using TOBREX while you are pregnant.
- are breast-feeding. TOBREX is not recommended during breastfeeding. It is unknown whether TOBREX will get into your milk.

Other warnings you should know about:

While you are using TOBREX

- **Driving and using machines:** your vision may be blurred for a time just after you apply TOBREX. Do not drive or use machines until this has worn off.
- **If you wear contact lenses:**
 - TOBREX Eye Drops contains a preservative (benzalkonium chloride) which may cause eye irritation and discolour soft contact lenses.
 - You should not wear contact lenses if you have an eye infection. If you are allowed to wear contact lenses, remove them before using TOBREX. Wait at least 15 minutes after using TOBREX to put your lenses back into your eyes.
- If the signs of your eye infection do not improve within 3 days or if they become worse, tell your healthcare professional.

Tell your healthcare professional about all the medicines you are using, have recently used or might use, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The use of TOBREX with other medicines has not been studied.

Tell your healthcare professional if you are using:

- Other antibiotics of the same family (aminoglycosides) such as amikacin, gentamycin, kanamycin, paromomycin, streptomycin

Always keep a list of your medicines and show it to your healthcare professional when you get a new medicine. It is important that your healthcare professional reviews all medications and supplements you are taking before prescribing TOBREX.

How to take TOBREX

Usual dose (adults and children 1 year and older):

The usual dose will depend on the type and severity of your infection


- In **mild to moderate eyes infections** your healthcare professional may recommend one drop or two drops into the affected eye(s), every four hours.
- In **severe eyes infections** this may be increased to two drops every hour.

Your healthcare professional will tell you how much you need to use each day. Ask your healthcare professional if you have any questions about the drug.

How to use:



1



2

1. Wash your hands.
2. Get the TOBREX bottle and a mirror.
3. **Shake well before use.**
4. After removing the cap: if the security snap collar is loose, remove it before using TOBREX.
5. Hold the bottle, pointing down, between your thumb and fingers.
6. Tilt your head back.
7. Pull down your lower eyelid with a clean finger until there is a “v” pocket between your eyelid and your eye. The drop will go in here (picture 1).
8. Bring the bottle tip close to the eye. Do this in front of a mirror if it helps.
9. **Do not touch your eye, eyelid, surrounding areas or other surfaces with the dropper, to avoid contaminating the suspension.**
10. Gently press on the base of the bottle to release one drop at a time. Do not squeeze the bottle. It is designed so that a gentle press on the bottom is all that it needs (picture 2).
11. If you miss, wipe up and try again.
12. Close the bottle immediately after use.

If you are using other eye drop or eye ointment medicines:

- Leave at least 5 minutes between each medicine.
- Instill all your eye drops before applying your eye ointment.

Overdose:

If you think you have taken too much or accidentally swallowed TOBREX, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you forget to use TOBREX, continue with the next dose as planned. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **Do not** use a double dose to make up for forgotten individual doses.

What are possible side effects from using TOBREX?

These are not all the possible side effects you may feel when taking TOBREX. If you experience any side effects not listed here, contact your healthcare professional. Please also see Warnings and Precautions.

The following side effects are common and about 3 in 100 people experienced them:

- itchy and swollen eyelid
- eye redness
- watery eye(s).

Other side effects include:

- eye problems such as:
 - eye discharge
 - dry, itchy or swollen eyes
 - increased tearing
 - corneal damage (the corneal is the clear surface of your eye)
 - blurred vision
 - eye allergy
 - loss of eyelashes
- headache
- skin problems such as swollen skin, hives, itchy or dry skin, white patches on the skin.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNKNOWN <ul style="list-style-type: none"> • Signs of an allergic reaction: • rash • hives • itching • red, swollen, blistered, or peeling skin with or without fever • wheezing • tightness in the chest or throat • trouble breathing or talking • unusual hoarseness • swelling of the mouth, face, lips, tongue, or throat 			✓
UNKNOWN Keratitis (an inflammation of the cornea — the clear front portion of your eye): <ul style="list-style-type: none"> • change in eyesight, • eye pain • sensitivity to light (photophobia) • difficulty opening your eyelid because of pain or irritation 			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.hc-sc.gc.ca/dhp-mps/medeff/report-declaration/index-eng.php>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store at room temperature (15°C - 30°C).
- Discard 28 days after initial opening. The best way to dispose of your medication is through a medicine take-back program. Check with your pharmacist about how to throw out unused medicines.
- Keep tightly closed.
- Keep out of the reach and sight of children.

If you want more information about TOBEX:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the [Health Canada website](#); the manufacturer's website www.novartis.ca, or by calling 1-800-363-8883.

This leaflet was prepared by Novartis Pharmaceuticals Canada Inc.

Last Revised: October 11, 2017

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

Pr **TOBREX® 0.3%**

Tobramycin Ophthalmic Ointment

Read this carefully before you start taking **TOBREX®** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **TOBREX**.

What is TOBREX used for?

TOBREX is used in adults and children above the age of 1 year to treat bacterial infections of the external part of the eye, such as conjunctivitis.

Antibacterial drugs like TOBREX treat only bacterial infections. They do not treat viral infections. Although you may feel better early in treatment, TOBREX should be used exactly as directed. Misuse or overuse of TOBREX could lead to the growth of bacteria that will not be killed by TOBREX (resistance). This means that TOBREX may not work for you in the future. Do not share your medicine.

How does TOBREX work?

TOBREX contains a medicine called tobramycin. Tobramycin belongs to the aminoglycoside class of antibiotic. Tobramycin works by killing bacteria that cause infections.

The signs of your eye infections [eye redness, increased tearing, discharge (pus or mucus) etc.] may start to improve within 3 days after starting TOBREX.

What are the ingredients in TOBREX?

Medicinal ingredients: tobramycin 0.3% w/w (3.0 mg/g)

Non-medicinal ingredients:

Preservative: chlorobutanol 0.5% w/w

Other ingredients: mineral oil, petrolatum base.

TOBREX comes in the following dosage forms:

TOBREX comes as eye drop and eye ointment.

Do not use TOBREX if you are allergic to:

- tobramycin,
- any of the other ingredients in TOBREX (see **What are the ingredients in TOBREX?**),
- other aminoglycoside antibiotics such as amikacin, gentamycin, kanamycin, paromomycin, streptomycin.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take TOBREX. Talk about any health conditions or problems you may have, including if you:

- are pregnant, you think you may be pregnant or are planning to have a baby. You will need to talk to your healthcare professional about the benefits and risks of using TOBREX while you are pregnant.
- are breast-feeding. TOBREX is not recommended during breastfeeding. It is unknown whether TOBREX will get into your milk.

Other warnings you should know about:

While you are using TOBREX

- **Driving and using machines:** your vision may be blurred for a time just after you apply TOBREX. Do not drive or use machines until this has worn off.
- **If you wear contact lenses:**
 - You should not wear contact lenses if you have an eye infection..
- If the signs of your eye infection do not improve within 3 days or if they become worse, tell your healthcare professional.

Tell your healthcare professional about all the medicines you use, have recently used or might use, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The use of TOBREX with other medicines has not been studied.

Tell your healthcare professional if you are using:

- Other antibiotics of the same family (aminoglycosides) such as amikacin, gentamycin, kanamycin, paromomycin, streptomycin

Always keep a list of your medicines and show it to your healthcare professional when you get a new medicine. It is important that your healthcare professional reviews all medications and supplements you are taking before prescribing TOBREX.

How to take TOBREX

Usual dose (adults and children 1 year and older):

The usual dose will depend on the type and severity of your infection

- In **mild to moderate eyes infections** your healthcare professional may recommend that you apply 1.25 cm (half inch) ribbon into the affected eye(s) two to three times per day.
- In **severe eyes infections** this may be increased to 1.25 cm (half-inch) ribbon into the affected eye(s) every three to four hours.

Your healthcare professional will tell you how much you need to use each day. Ask your healthcare professional if you have any questions about the drug.

How to use:



1. Wash your hands.
2. Tilt your head back.
3. Place a finger on your cheek just under your eye and gently pull down until a “v” pocket is formed between your eyeball and lower eyelid.
4. Place a small amount (1.25 cm or half inch ribbon) of TOBREX in the “v” pocket. **Do not let the tip of the tube touch your eye**, to avoid contaminating the ointment.
5. Look down before closing your eye.
6. Replace the cap of the tube.

If you are using other eye drop or eye ointment medicines:

- Leave at least 5 minutes between each medicine.
- Instill all your eye drops before applying your eye ointment.

Overdose:

If you think you have taken too much or accidentally swallowed TOBREX, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you forget to use TOBREX, continue with the next dose as planned. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. **Do not** use a double dose to make up for forgotten individual doses.

What are possible side effects from using TOBREX?

These are not all the possible side effects you may feel when taking TOBREX. If you experience any side effects not listed here, contact your healthcare professional. Please also see Warnings and Precautions.

The following side effects are common and about 3 in 100 people experienced them:

- itchy and swollen eyelid
- eye redness
- watery eye(s).

Other side effects include:

- eye problems such as:
 - eye discharge
 - dry, itchy or swollen eyes
 - increased tearing
 - corneal damage (the corneal is the clear surface of your eye)
 - blurred vision
 - eye allergy
 - loss of eyelashes
- headache
- skin problems such as swollen skin, hives, itchy or dry skin, white patches on the skin.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<p>UNKNOWN</p> <ul style="list-style-type: none"> • Signs of an allergic reaction: • rash • hives • itching • red, swollen, blistered, or peeling skin with or without fever • wheezing • tightness in the chest or throat • trouble breathing or talking • unusual hoarseness • swelling of the mouth, face, lips, tongue, or throat 			✓
<p>UNKNOWN</p> <p>Keratitis (an inflammation of the cornea — the clear front portion of your eye):</p> <ul style="list-style-type: none"> • change in eyesight, • eye pain • sensitivity to light (photophobia) • difficulty opening your eyelid because of pain or irritation 			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.hc-sc.gc.ca/dhp-mps/medeff/report-declaration/index-eng.php>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store at room temperature (15°C - 30°C).
- Discard 28 days after initial opening. The best way to dispose of your medication is through a medicine take-back program. Check with your pharmacist about how to throw out unused medicines.
- Keep tightly closed.
- Keep out of the reach and sight of children.

If you want more information about TOBREX:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the [Health Canada website](#); the manufacturer's website www.novartis.ca, or by calling 1-800-363-8883.

This leaflet was prepared by Novartis Pharmaceuticals Canada Inc.

Last Revised: October 11, 2017